

**IMPORTANT: PLEASE CONTINUE TAKING ALL BLOOD PRESSURE MEDICATION ON THE DAY OF PROCEDURE.**

### **MEDICATIONS TO BE STOPPED PRIOR TO PROCEDURE**

#### **Blood Thinners:**

**HOLD FOR 2 DAYS (48 HOURS):** Xarelto or Eliquis

**HOLD FOR 5 DAYS:** Plavix, Brilinta, Coumadin, or Pradaxa

#### **Diabetic Medications/also used for weight loss:**

**HOLD FOR 2 DAYS (48 HOURS):** Metformin, Byetta, Saxenda and Victoza

**HOLD FOR 2 WEEKS (14 DAYS):** Trulicity (Dulaglutide), Mounjaro (Tirzepatide)  
Ozempic/Wegovy/Rybelsus (Semaglutide)

**INSULIN:** Take only ½ the regular PM dose of insulin the evening prior to the procedure. Check your glucose level the morning of your procedure. Hold insulin if the level is low, otherwise you may take half the regular insulin in the morning.

#### **OTHER MEDICATION:**

**HOLD FOR 7 DAYS:** Phentermine

**HOLD ON THE DAY OF PROCEDURE:** Furosemide (Lasix)

**HOLD FOR 48 HOURS:** MARIJUANA

ANY QUESTIONS OR CONCERNS REGARDING MEDICATIONS, PLEASE CALL THE OFFICE AND SPEAK TO A NURSE 520-795-3090. FAILURE TO COMPLY CAN RESULT IN CANCELLATION OF PROCEDURE.

**YOU DO NOT NEED TO STOP NSAIDS PRIOR TO THE PROCEDURE (ALEVE, IBUPROPHEN, ASPIRIN ECT)**

